

Population Dynamics of the Root-knot Nematodes *Meloidogyne incognita* (Kofoed & White) Chitwood and *M. javanica* (Treub) Chitwood on Grapevines in two different Regions of South Africa.¹

J.T. LOUBSER (a) and A.J. MEYER (b)

a. Viticultural and Oenological Research Institute, Private Bag X5026, 7600 Stellenbosch, Republic of South Africa.

b. Department of Entomology and Nematology, University of Stellenbosch, 7600 Stellenbosch, Republic of South Africa.

Submitted for publication: June 1987

Accepted for publication: September 1987

Keywords: Population dynamics, root-knot nematodes, *Meloidogyne incognita*, *Meloidogyne javanica*, grapevines.

Two root-knot nematode species, *Meloidogyne incognita* and *M. javanica*, were studied with regard to their seasonal population fluctuations on grapevines growing in two vastly different climatic areas. Regular observations on reproduction and numbers of larvae in the soil were compared with patterns of root growth, soil temperature and moisture.

Population fluctuations of the two species showed similar trends in spite of the climatic differences in the two areas, but *M. incognita* in the northern Cape reached higher populations. Larvae populations in the soil declined in summer in both areas and increased during autumn to reach peaks in winter. With the onset of root growth in spring, larvae numbers decreased in the soil, as a result of large scale root penetration.

Present knowledge of root-knot nematode distribution in the root area of grapevines and the seasonal population fluctuations is largely inadequate. Root-knot nematode population dynamics is important because it forms the basis for advisory work about these nematodes. This lack of information may also have contributed to the unsatisfactory chemical control achieved in established vineyards in South Africa (Loubser & De Klerk, 1986) as well as in other countries (Raski *et al.*, 1981; Harris, 1986). No study has yet been made of the bionomics of *Meloidogyne* species in South African vineyards.

Temperature, humidity, light, aeration of the soil, age and nutritional status of the host may influence the biological activities in the life cycle and development of *Meloidogyne* spp. (De Guiran & Ritter, 1979) while both the host plant and its environment will influence the population dynamics of these parasites (Ferris & Van Gundy, 1979). Root-knot nematode populations are therefore thought to fluctuate between soil types, different hosts and different geographical locations. However, it was shown to follow the same pattern every year both on monocultured annuals (Johnson, Dowler & Hauser, 1974) and perennial crops such as grapevines (Ferris & McKenry, 1976a). Therefore, although soil temperature and soil moisture play an important role in nematode population numbers on grapevine as reported by Ferris & McKenry (1974, 1976b), these may only be rate modifying factors which will not influence the nature of the annual population curve on a specific host plant.

The present study was carried out in order to learn more of the bionomics of *Meloidogyne incognita* and *M. javanica* on grapevines under two different climatic conditions in South Africa.

MATERIALS AND METHODS

Experimental vineyards

Two experimental plots were used: 1) A flood-irrigated vineyard on a loamy sand (Table 1) in the summer rainfall area (Vaalharts, Northern Cape Province) with a high infestation of *Meloidogyne incognita*. 2) A microjet-irrigated vineyard on sandy loam (Table 1) in the winter rainfall region (Bien Donné, Western Cape Province) infested with *Meloidogyne javanica*. Both vineyards were approximately 12 years old with Jacques as rootstock.

TABLE 1

Soil characteristics of trial vineyards in Vaalharts and Bien Donné.

Depth (mm)	Location	Sand (%)			Silt (%)	Clay (%)	pH (KCl)	R (ohms)
		Fine	Medium	Coarse				
300	Vaalharts	73.9	17.0	1.0	1.1	7.0	5.7	1700
	Bien Donné	45.3	21.6	3.7	16.6	12.8	4.4	2900
600	Vaalharts	74.3	16.1	0.9	0.8	7.9	5.6	1800
	Bien Donné	42.2	6.4	1.6	30.0	19.8	4.2	4400
900	Vaalharts	71.4	16.3	0.9	1.2	10.2	5.4	2200
	Bien Donné	33.0	11.8	10.6	29.6	15.0	4.1	4500

Sampling procedures

a. Soil: The Vaalharts vineyard consisted of four adjacent blocks of 25 vines each, planted in five rows (3,75 m apart) with five vines (1,5 m apart) per row. The experimental area for the Bien Donné vineyard consisted of ten rows of 40 vines each, with a vine spacing of 3,3 m x 1,83 m.

¹ Part of a thesis to be submitted by the senior author to the University of Stellenbosch for the Ph. D (Agric.) degree.

An initial survey was conducted to determine the best sampling procedure for optimum nematode recovery in both plots because of the differences in layout. Based on these results, the four vineyard blocks of the Vaalharts vineyard were sampled separately, each by bulking 25 soil cores taken in the root zone (10-450 mm depth) with a 25 mm \emptyset auger within 150-450 mm from the trunks of individual vines. The Bien Donn  vineyard was sampled in the same manner but samples consisted of 40 soil cores each.

Because of the low number of vines in the Vaalharts trial and because sampling was performed on a weekly basis, soil cores had to be taken from the same vines every week. To avoid concentration of the root damage, consecutive samples were taken in a clockwise direction and towards the vine trunk after completing a full circle. At Bien Donn , sampling was done randomly every two weeks.

b. Roots: In order to establish periods of maximum nematode reproduction, root samples were collected weekly from four separate vines (one per block) and a 30 g aliquot was used for extracting eggs and larvae from each sample. This survey was conducted at Vaalharts only.

Extraction procedures

Samples were placed in plastic bags and processed 1-4 hours after collection. Analyses for second stage larvae in the soil were done by a motility-independent sieving-sedimentation method (Loubser, 1985) while root analyses for eggs and larvae were done according to the method of Hussey & Barker (1973). For each egg suspension the embryonic development was recorded by distinguishing between undifferentiated eggs and eggs developed to the first larval stage. Second stage larvae in the egg sacs were also recorded.

Root growth

Root growth was measured only in the Vaalharts vineyard. This was done by means of underground observation chambers which allowed observation of four vines. Details of this study are discussed by Loubser & Meyer (1986).

Soil temperatures and soil moisture

Soil temperatures were measured at 150 mm, 300 mm and 600 mm depths on a three-hourly basis by means of soil thermistors coupled to a micrologger. Soil moisture was measured at 300 mm and 600 mm depth every second or third day by means of a mercury tensiometer. The average monthly soil temperature and soil moisture were calculated for both trial plots as well as the total monthly heat units (in degree days) as described by Tyler (1933).

RESULTS AND DISCUSSION

Vaalharts trial

Soil population fluctuations of *Meloidogyne incognita* are shown as the average monthly number of second-stage larvae recorded (Fig. 1 A). Populations were low in summer (January), and apart from a drop during June, increased gradually to reach a peak in midwinter (July). Thereafter populations declined gradually to reach another low during summer the next season (December). Observations during the second year in the same vineyard confirmed these results. Similar results

were obtained by Ferris & McKenry (1974) in California.




FIG. 1

Average monthly figures for number of *Meloidogyne incognita* larvae in the soil (A), reproduction as measured by eggs and larvae per 1 g of roots (B), soil temperature (C), soil moisture (D) and root growth (E) of a vineyard at Vaalharts.

Soil temperatures at 150 mm, 300 mm and 600 mm differed very little and fluctuated slightly on a daily basis. For this reason only the average monthly temperature at 300 mm depth is shown in Fig. 1 C. Soil populations of the nematode reached a peak when soil temperatures were at their lowest (12°C) and were low when soil temperatures were higher (max. 29°C). During late winter most larvae were found in a coiled position, presumably responding to environmental stress (eg. low soil temperatures) by quiescence (De Guiran, 1979a).

The average monthly soil moisture levels at 300 mm and 600 mm depth are shown in Fig. 1 D. Soil moisture was relatively constant over most of the sampling period except for a steep drop during time of ripening of grapes (February and March). Soil larval populations were thus not markedly influenced by soil moisture.

The average monthly root growth is presented in Fig. 1 E. It shows a high number of new root tips in March and again during October, November and December. Very little root growth was recorded during June, July

and August. At this stage larval populations in the soil were at their highest, probably because of the lack of available infestation sites and because of low activity due to low soil temperatures. The increase in numbers of larvae in the soil preceding this period can be explained by the decrease in new root growth as well as the reproduction of established females. At the onset of new root growth in spring (September), larval populations in the soil decreased rapidly to reach a low which persisted during summer.

According to Ferris & McKenry (1974) root infestation in spring is primarily the result of newly hatched larvae because overwintering larvae appear to be of low infectivity. Although these larvae were found to be vacuolated as reported by the above workers, they only became so as soil temperatures increased during early spring and they became active. We believe that overwintering larvae constituted a high proportion of the invading population in the Vaalharts vineyards. Since the motility-independent extraction method which was used did not show a large number of dead larvae in the soil, these larvae must have penetrated newly developed roots.

Reproduction as expressed by the monthly average number of eggs and larvae extracted from roots, is shown in Fig. 1 B. The curve shows two definite peaks, one in late summer (February) and another during spring (September). Reproduction apparently fluctuated regardless of soil temperatures since both increases and decreases occurred with rising temperatures. On the other hand, nematode reproduction did remain relatively constant during winter, i.e. larvae and eggs were always present. The latter may represent eggs in diapause (De Guiran, 1979b) or quiescence (Linford, 1941). The fact that larval populations in the soil increased during winter, suggests, however, that reproduction or at least hatching continued throughout winter.

When reproduction is compared with soil moisture, no connection is found. A reproduction peak was observed during February when soil moisture was at its lowest and again during September when soil moisture was high. Soil moisture never dropped below 4-5%, the level at which egg hatch is affected (Ferris & McKenry, 1974).

It is known that soil temperature and soil moisture play an important role in the root-knot nematode's biology (Wallace, 1963). The effect they had on reproduction and development in this study was probably masked by the combined influence of all factors involved. Increasing soil moisture after harvest in March could have stimulated egg hatch and given rise to more second-stage larvae in the soil. This could have led to an increase in root infestation at a stage when new root growth occurred. The second reproduction peak in spring (September) was probably triggered by rising soil temperatures and coincided with another root growth flush.

Embryonic development and eclosion (i.e. the escape of the larvae through the egg shell) of root-knot nematode eggs in the Vaalharts vineyard, can be followed from the results listed in Table 2. The percentage undifferentiated eggs, which may represent quiescence (Linford, 1941) or diapause (De Guiran, 1979b), did

not increase during the observation period, suggesting that they did not occur. Wallace (1971) found in glass-house experiments that eclosion, but not embryonic development, is arrested at low temperatures. This would have led to lower numbers of second stage larvae in the egg mass and a higher percentage of developed eggs during winter. At higher temperatures, on the other hand, embryonic development is inhibited (Wallace, 1971). The field results of the present study do not substantiate either of these two observations. The percentage of developed eggs decreased rather than increased during winter. This emphasizes the enormous gap between results sometimes obtained under controlled conditions and field trials.

TABLE 2

Seasonal egg development of *Meloidogyne incognita* on grapevine roots in Vaalharts vineyards.¹

Month	Undifferentiated eggs	Developed eggs (J ₁ -stage)	Larvae (J ₂ -stage)
January	54	10	36
February	53	16	31
March	52	11	37
April	41	9	50
May	55	8	37
June	54	9	37
July	42	8	50
August	59	6	35
September	51	17	32
October	53	16	31
November	63	16	21
December	67	7	26

1. Each developmental stage is expressed as a percentage of the total population extracted from 120 g roots (4 x 30 g replicates).

Bien Donné trial

Fluctuations in numbers of *Meloidogyne javanica* larvae in the soil, soil temperature and soil moisture are shown in Fig. 2. Larval population trends are essentially similar to those of *M. incognita* in the Vaalharts trial with low numbers occurring in summer and high numbers occurring in winter. Much lower numbers of *M. javanica* larvae were present in the soil compared to the Vaalharts vineyard. This may be attributed to species differences, but ecological factors should also be considered.

Both surveys were conducted on Jacquez rootstock, but the soils involved were different (Table 1) and could have influenced nematode activities. Temperatures varied in a similar manner but with a mean minimum of 11°C in winter and a mean maximum of 22°C in summer. Soil moisture levels fluctuated between 7% and 14% as compared to 6% and 13% for the Vaalharts trial. Apart from a drop in soil moisture in May, the average soil moisture was ca. 11%; almost the same as for the Vaalharts soil. According to Ferris, Schneider & Semenov (1984), temperature is the major extrinsic reproduction rate-determining factor and this is most probably responsible for the differences recorded between populations in the two vineyards. The number of degree days (DD₁₀) for both trials, are listed in Table 3




FIG. 2.

Average monthly figures for number of *Meloidogyne javanica* larvae in the soil (A), soil temperature (B), and soil moisture (C) in a vineyard at Bien Donné.

TABLE 3

Degree days (10°C) calculated for trial vineyards at Vaalharts and Bien Donné.

Month	Degree days (DD_{10})	
	Vaalharts	Bien Donné
January	588	376
February	510	326
March	567	330
April	316	183
May	263	249
June	115	135
July	78	84
August	151	121
September	218	159
October	399	273
November	386	280
December	445	314
Total:	4036	2830

for each month as well as accumulatively for the full sampling period.

Heat-units (Tyler, 1933) have been used to relate soil temperatures to *Meloidogyne arenaria* penetration and development in grapevines (Ferris & Hunt, 1979; Ferris, Schneider & Stuth, 1982; Ferris, Schneider & Semenoff, 1984). The number of DD_{10} required by this nematode to reach maturity, differed between grapevine cultivars. From the results of Ferris & Hunt (1979), 667 DD_{10} was assumed to be necessary for any *Meloidogyne* species to develop from the egg stage to

maturity on a susceptible rootstock such as Jacquez. Based on this assumption, it was calculated that the number of root-knot nematode generations will be 6.05 and 4.24 per annum in the Vaalharts and Bien Donné vineyards respectively. These relative figures for numbers of generations in the two species examined may partially explain the differences recorded in nematode numbers in the different trial plots. It remains, however, necessary to investigate the reproductive potential of the two *Meloidogyne* species involved in more detail in order to determine its role in explaining population numbers.

CONCLUSIONS

In spite of differing climatic and other ecological conditions, *M. incognita* and *M. javanica* showed a cyclical fluctuation in numbers that was largely similar on the same host plant. The differences that were detected consisted merely of differences in the magnitude of the populations which was possibly the result of the greater number of generations on the site with the higher soil temperatures.

The results support findings by other workers with regard to seasonal fluctuations of root-knot nematodes. Although strongly affected by soil and climatic conditions, the population dynamics of obligate parasites such as root-knot nematodes, seems closely related to the host plant, especially with regard to root growth periods. This fact should be considered in any pest management programme in order to achieve improved nematode control.

Quiescence and diapause were not observed in the present study. They either did not occur in the vineyards under observation or the techniques used were not sensitive enough to detect them.

The information obtained in this study on the population dynamics of the two *Meloidogyne* species, coupled with our knowledge of grapevine root distribution and nematicide persistence in the soil, provides a solid theoretical basis for recommendations on root-knot nematode control. Appropriately, chemical control in established vineyards should commence immediately after harvest and/or during early spring. During these stages new root growth is initiated which should be protected against infestation by second stage larvae. However, the relative importance of infection of the two root growth flushes, is still unknown. Furthermore, importance of early applications, prior to infestation, should also be determined in order to establish the most effective time of application.

LITERATURE CITED

- DE GUIRAN, G., 1979a. Survie des nématodes dans les sols secs et saturés d'eau: oeufs et larves de *Meloidogyne incognita*. *Revue Nematol.* **2**, 65-77.
- DE GUIRAN, G., 1979b. A necessary diapause in root-knot nematodes. Observations on its distribution and inheritance in *Meloidogyne incognita*. *Revue Nematol.* **2**, 223-231.
- DE GUIRAN, G. & RITTER, M., 1979. Life cycle of *Meloidogyne* species and factors influencing their development. In: F. Lamberti and C.E. Taylor (eds.) Root-knot nematodes (*Meloidogyne* species); systematics, biology and control. Acad. Press, London, pp. 173-191.
- FERRIS, H. & HUNT, W.A., 1979. Quantitative aspects of the development of *Meloidogyne arenaria* larvae in grapevine varieties and rootstocks. *J. Nematol.* **11**, 168-174.

- FERRIS, H. & MCKENRY, M., 1974. Seasonal fluctuations in the spatial distribution of nematode populations in a California vineyard. *J. Nematol.* **6**, 203-210.
- FERRIS, H., & MCKENRY, M., 1976a. A survey of nematode distribution in California vineyard soils. *J. Amer. Soc. Hort. Sci.* **161**, 332-336.
- FERRIS, H. & MCKENRY, M., 1976b. Nematode community structure in a vineyard soil. *J. Nematol.* **8**, 131-137.
- FERRIS, H., SCHNEIDER, S.M. & SEMENOFF, M.C., 1984. Distributed egg production functions for *Meloidogyne arenaria* in grape varieties and consideration of the mechanistic relationship between plant and parasite. *J. Nematol.* **16**, 178-183.
- FERRIS, H., SCHNEIDER, S.M. & STUTH, M.C., 1982. Probability of penetration and infection by root-knot nematode, *Meloidogyne arenaria*, in grape cultivars. *Am. J. Enol. Vitic.* **33**, 31-35.
- FERRIS, H. & VAN GUNDY, S.D., 1979. *Meloidogyne* ecology and host interrelationships. In: F. Lamberti and C.E. Taylor (eds.) Root-knot nematodes (*Meloidogyne* species); systematics, biology and control. Acad. Press, London, pp. 205-230.
- HARRIS, A.R., 1986. Comparison of some nematicides on *Vitis vinifera* cv. Sultana in Victoria, Australia. *Am. J. Enol. Vitic.* **37**, 224-227.
- HUSSEY, R.S. & BARKER, K.R., 1973. A comparison of methods of collecting inocula of *Meloidogyne* spp., including a new technique. *Pl. Dis. Reprtr* **57**, 1025-1028.
- JOHNSON, A.W., DOWLER, C.C. & HAUSER, E.W., 1974. Seasonal population dynamics of selected plant-parasitic nematodes on four monocultured crops. *J. Nematol.* **6**, 187-190.
- LINFORD, M.B., 1941. Some soil moisture relationships of the root-knot nematode. *Phytopathology* **31**, 862.
- LOUBSER, J.T., 1985. A modified sieving-sedimentation method for extracting nematodes from soil. *Hort. Sci.* **3**, 23-25.
- LOUBSER, J.T. & DE KLERK, C.A., 1986. Chemical control of root-knot nematodes in established vineyards. *S. Afr. J. Enol. Vitic.* **6**, 31-33.
- LOUBSER, J.T. & MEYER, A.J., 1986. Strategies for chemical control of root-knot nematodes (*Meloidogyne incognita*) in established vineyards. *S. Afr. J. Enol. Vitic.* **7**, 84-89.
- RASKI, D.J., JONES, N.O., HAFEZ, S.L., KISSLER, J.J. & LUVISKI, D.A., 1981. Systemic nematicides tested as alternatives to DBCP. California Agriculture, May-June, 11-12.
- TYLER, J., 1933. Development of the root-knot nematode as affected by temperature. *Hilgardia* **7**, 391-415.
- WALLACE, H.R., 1963. The biology of plant parasitic nematodes. Edward Arnold (Publishers) Ltd., London.
- WALLACE, H.R., 1971. The influence of temperature on the embryonic development and hatch in *Meloidogyne javanica*. *Nematologica* **17**, 179-186.