

Errata

Myburgh, P.A., 2003. Responses of *Vitis vinifera* L. cv. Sultanina to water deficits during various pre- and post-harvest phases under semi-arid conditions. S. Afr. J. Enol. Vitic. 24, 25-33.

Table 4 on page 30 must be replaced by the following:

TABLE 4

Phosphorus and cation concentrations as well as Ca:Na ratios in flesh and skins of yellow and brown dipped Sultanina raisins as measured during the 1996/97 season.

Treatment number	Raisin colour	Flesh/skins	P (%)	Ca (%)	Mg (%)	K (%)	Na (%)	Ca:Na
T1	Yellow	Flesh	0.49	0.52	0.16	0.83	0.014	37.1
		Skins	0.45	0.51	0.15	0.82	0.017	30.0
	Brown	Flesh	0.46	0.44	0.18	0.98	0.027	16.3
		Skins	0.36	0.53	0.14	1.39	0.034	15.6
T5	Yellow	Flesh	0.35	0.31	0.13	1.00	0.012	25.8
		Skins	0.44	0.54	0.14	0.98	0.015	36.0
	Brown	Flesh	0.40	0.31	0.15	1.13	0.032	9.7
		Skins	0.47	0.44	0.18	1.11	0.043	10.2
LSD ($p \leq 0.05$)			NS*	NS	NS	NS	0.028	NS

* NS = Not significant.